

**M E T U Department of Mathematics**

<b>Math 111 Fundamentals of Mathematics Fall 2018 Final Exam 15.01.2019 13:30</b>									
Last Name :				Section :					
Name :				Duration : 120 <i>minutes</i>					
Student No :									
7 QUESTIONS ON 4 PAGES							TOTAL 80 POINTS		
1	2	3	4	5	6	7			

By signing below, I pledge that I will write this examination as my own work and without the assistance of others or the usage of unauthorized material or information. I understand that possession of any kind of electronic device during the exam is prohibited. I also understand that not obeying the rules of the examination will result in immediate cancellation and disciplinary procedures.

Signature .....

**(5+5+5 pts) 1.** The parts of this question are unrelated.

- a) Let  $A, B$  be sets such that  $A \subseteq B$ . Show that if  $B$  is uncountable and  $A$  is countable, then  $B - A$  is uncountable.

Assume towards a contradiction that  $B$  is uncountable and  $A$  is countable, and  $B - A$  is countable. Since  $A \subseteq B$ , we have that  $B = A \cup (B - A)$ . Because the union of two countable sets is countable and,  $A$  and  $B - A$  are both countable, we have that  $B$  is countable, which is a contradiction.

- b) Show that the following statement is false:


For all sets  $A$  and  $B$ , if  $A$  and  $B$  are uncountable, then  $A \cap B$  is uncountable.

We shall give a counterexample to this statement. Choose  $A = (0, 1)$  and  $B = (-1, 0)$ . Then, we have that  $A \sim B \sim \mathbb{R}$  and hence both  $A$  and  $B$  are uncountable. On the other hand,  $A \cap B = \emptyset$ , which is countable.

- c) Prove that for any set  $A$  and  $B$ , we have that  $\mathcal{P}(A) \cap \mathcal{P}(B) \subseteq \mathcal{P}(A \cap B)$ .

Let  $A$  and  $B$  be sets. We will prove that  $\mathcal{P}(A) \cap \mathcal{P}(B) \subseteq \mathcal{P}(A \cap B)$ .

Let  $S \in \mathcal{P}(A) \cap \mathcal{P}(B)$ . Then we have that  $S \in \mathcal{P}(A)$  and  $S \in \mathcal{P}(B)$ , and so  $S \subseteq A$  and  $S \subseteq B$ . It follows from  $S \subseteq A$  and  $S \subseteq B$  that  $S \subseteq A \cap B$ , and hence  $S \in \mathcal{P}(A \cap B)$ .



**(5+5+5 pts) 2.** Let  $X = \{0, 1, 2, 3, 4, \dots\}$ ,  $[0, \infty) = \{x \in \mathbb{R} : 0 \leq x\}$  and  $(0, 1) = \{x \in \mathbb{R} : 0 < x < 1\}$ . Consider the function

$$f : X \times (0, 1) \rightarrow [0, \infty)$$

defined by  $f(n, r) = n + r$  for all  $(n, r) \in X \times (0, 1)$ .

a) Show that the function  $f$  is injective.

Let  $(n, r)$  and  $(m, s)$  be elements of  $X \times (0, 1)$ . Assume that  $f(n, r) = f(m, s)$ . Then we have that  $n + r = m + s$  and so  $n - m = s - r$ . Since  $m$  and  $n$  are integers,  $n - m$  is an integer. Since  $0 < r, s < 1$ , we have that  $-1 < s - r = n - m < 1$ . This implies that  $n - m = 0$  and hence  $n = m$ . Consequently,  $s = r$  and so  $(n, r) = (m, s)$ . This shows that  $f$  is injective.

b) Show that the function  $f$  is not surjective.

Notice that  $0 \in [0, \infty)$ . We claim that there exists no  $(n, r) \in X \times (0, 1)$  such that  $f(n, r) = 0$  which will show that  $f$  is not surjective. Let  $(n, r) \in X \times (0, 1)$ . Then  $f(n, r) = n + r > 0 = 0$ . Hence  $f(n, r) \neq 0$ .

c) You are **given** the fact that there exists an injective function  $g : [0, \infty) \rightarrow X \times (0, 1)$ . Show that  $X \times (0, 1)$  and  $[0, \infty)$  have the same cardinality.

We have proven in (a) that there exists an injection  $f : X \times (0, 1) \rightarrow [0, \infty)$  and we are given that there exists an injection  $g : [0, \infty) \rightarrow X \times (0, 1)$ . By Cantor-Schröder-Bernstein theorem, there exists a bijection between  $X \times (0, 1)$  and  $[0, \infty)$  which shows that these sets have the same cardinality.

**(5 pts) 3.** Let  $A, B, C, D$  be sets and  $f : A \rightarrow B$  and  $g : C \rightarrow D$  be functions. Consider the function  $h : A \times C \rightarrow B \times D$  given by  $h(a, c) = (f(a), g(c))$  for all  $(a, c) \in A \times C$ . Show that if  $f$  and  $g$  are surjective, then  $h$  is surjective.

Assume that  $f$  and  $g$  are surjective. Let  $(b, d) \in B \times D$ . Then  $b \in B$  and  $d \in D$ . Since  $f$  and  $g$  are surjective, there exist  $a \in A$  and  $c \in C$  such that  $f(a) = b$  and  $g(c) = d$ . Note that  $(a, c) \in A \times C$  and  $h(a, c) = (f(a), g(c)) = (b, d)$ . Thus  $h$  is surjective.

(5 pts) 4. Using **induction**, prove that

$$1 + 3 + 5 + \dots + (2n - 1) = n^2 \text{ for all natural numbers } n \geq 2.$$

**Base case.** We have that  $1 + (2 \cdot 2 - 1) = 1 + 3 = 4 = 2^2$ , and hence the claim holds for  $n = 2$ .

**Inductive step.** Let  $n \geq 2$  be an integer. Assume that  $1 + 3 + 5 + \dots + (2n - 1) = n^2$ . Then, by adding  $2n + 1$  to both sides, we have that

$$1 + 3 + 5 + \dots + (2n - 1) + (2n + 1) = n^2 + (2n + 1)$$

$$1 + 3 + 5 + \dots + (2n - 1) + (2(n + 1) - 1) = (n + 1)^2$$

and hence the claim holds for  $n + 1$ . It follows from the principle of induction that  $1 + 3 + 5 + \dots + (2n - 1) = n^2$  for all natural numbers  $n \geq 2$ .

(10+5 pts) 5. Let  $\mathbb{N} = \{1, 2, 3, \dots\}$  be the set of natural numbers. Consider the relation  $\sim$  on  $\mathbb{N}$  defined by

$$x \sim y \quad \text{if and only if} \quad \frac{y}{x} = 2^i \quad \text{for some } i \in \mathbb{Z}$$

for all  $x, y \in \mathbb{N}$ .

a) Show that  $\sim$  is an equivalence relation.

Let  $x \in \mathbb{N}$ . Then we have that  $\frac{x}{x} = 2^0$  and  $0 \in \mathbb{Z}$ . Thus, by definition,  $x \sim x$ . Hence,  $\sim$  is reflexive.

Let  $x, y \in \mathbb{N}$ . Assume that  $x \sim y$ . Then, by definition,  $\frac{x}{y} = 2^i$  for some  $i \in \mathbb{Z}$ . This implies that  $\frac{y}{x} = \frac{1}{2^i} = 2^{-i}$  and hence  $y \sim x$ . Therefore,  $\sim$  is symmetric.

Let  $x, y, z \in \mathbb{N}$ . Assume that  $x \sim y$  and  $y \sim z$ . Then, by definition,  $\frac{x}{y} = 2^i$  and  $\frac{y}{z} = 2^j$  for some  $i, j \in \mathbb{Z}$ . It follows that  $\frac{x}{z} = \frac{x}{y} \cdot \frac{y}{z} = 2^i \cdot 2^j = 2^{i+j}$ . Hence,  $x \sim z$  and so  $\sim$  is transitive.

a) Construct a bijection from  $\mathbb{N}$  to  $[1]$ .

We have that


$$[1] = \{n \in \mathbb{N} : 1 \sim n\} = \{n \in \mathbb{N} : \exists i \in \mathbb{Z} \frac{n}{1} = 2^i\} = \{1, 2, 4, 8, \dots\}$$

Consider the function  $f : \mathbb{N} \rightarrow [1]$  given by  $f(i) = 2^{i-1}$  for all  $i \in \mathbb{N}$ . Then  $f$  is a bijection.

(5 pts) 6. Let  $X = \{2, 3, 4, 5, 6, 7, 8\}$ . Consider the relation  $\preceq$  on the set  $X$  defined by

$$x \preceq y \quad \text{if and only if} \quad \frac{y}{x} = 2^i \quad \text{for some } i \in \mathbb{N} \cup \{0\}$$

for all  $x, y \in X$ . You are **given** that  $\preceq$  is a partial order relation. Draw the Hasse diagram of  $\preceq$ .



(5+5+5+5 pts) 7. The following proofs are **incorrect**. Briefly explain the mistake in each of the following proofs.

**Theorem.** Let  $A, B, C$  be sets. If  $(A \cap B) \subseteq C$ , then  $(A - C) \cap (B - C) = \emptyset$ .

**Proof.** We shall prove this statement by proving the contrapositive. Assume that  $(A \cap B) \not\subseteq C$ . Then there exists  $a \in A \cap B$  such that  $a \notin C$ . Since  $a \in A \cap B$ , we have that  $a \in A$  and  $a \in B$ . It follows from  $a \in A$  and  $a \notin C$  that  $a \in A - C$ . Similarly, it follows from  $a \in B$  and  $a \notin C$  that  $a \in B - C$ . Therefore,  $a \in (A - C) \cap (B - C)$  and hence  $(A - C) \cap (B - C) \neq \emptyset$ .

**Mistake in the proof:** The mistake in the proof is that the argument does not prove the contrapositive of the given statement. The contrapositive of the statement is that if  $(A - C) \cap (B - C) \neq \emptyset$ , then  $(A \cap B) \not\subseteq C$ .

**Theorem.** For all  $i \in \mathbb{Z}$ , there exists  $j \in \mathbb{Z}$  such that for all  $k \in \mathbb{N}$  we have  $2^i \cdot 2^j \leq k + 1$ .

**Proof.** Let  $i \in \mathbb{Z}$  and  $k \in \mathbb{N}$ . Choose  $j = -k - i$ . Then  $j \in \mathbb{Z}$  and

$$2^i \cdot 2^j = 2^i \cdot 2^{-k-i} = 2^{i-k-i} = 2^{-k} = \frac{1}{2^k} \leq 1 \leq k + 1$$

**Mistake in the proof:** The mistake in the proof is that the integer  $j$  cannot depend on the natural number  $k$ . For each  $i \in \mathbb{Z}$ , one should find some  $j \in \mathbb{N}$  which works for **every**  $k \in \mathbb{N}$  at the same time.

**Theorem.** Let  $\sim$  be the relation on  $\mathbb{N} = \{1, 2, 3, \dots\}$  defined by

$$x \sim y \quad \text{if and only if} \quad \frac{y}{x} = 2^i \quad \text{for some } i \in \mathbb{Z}$$

for all  $x, y \in \mathbb{N}$ . The relation  $\sim$  is reflexive.

**Proof.** Let  $x \in \mathbb{N}$ . Assume that  $x \sim x$ . Then, we have that  $\frac{x}{x} = 1 = 2^0$  and  $0 \in \mathbb{Z}$ . Therefore, the relation is reflexive.

**Mistake in the proof:** The mistake in the proof is that the argument assumes the statement  $x \sim x$  at the beginning. That  $x \sim x$  should be the conclusion of the argument, not the assumption.

**Theorem.** For any set  $A$ , if  $\mathcal{P}(A)$  is countable, then  $A$  is countable.

**Proof.** Let  $A$  be a set. Assume that  $\mathcal{P}(A)$  is countable. Since  $A \subseteq \mathcal{P}(A)$  and every subset of a countable set is countable,  $A$  is countable.

**Mistake in the proof:** The mistake in the proof is that it is not true in general that  $A \subseteq \mathcal{P}(A)$ .